

Association between index venous thromboembolism (VTE) event and 12-months outcomes for patients in routine clinical practice enrolled in the Edoxaban Treatment in routine cliNical prActice for patients with acute VTE in Europe (ETNA-VTE-Europe) registry

Michiel Coppens¹, Alexander T. Cohen², Cihan Ay³, Philippe Hainaut⁴, Ulrich Hoffmann⁵, Sean Gaine⁶, David Jiménez⁷, Marc Schindewolf⁸, Bernd Brüggengjürgen⁹, Pierre Levy¹⁰, Julio Lopez Bastida¹¹, Eric Vicaut¹², Peter Bramlage¹³, Giancarlo Agnelli¹⁴ and on behalf of the ETNA-VTE-Europe investigators

¹Department of Vascular Medicine, Amsterdam University Medical Centers, University of Amsterdam, Amsterdam, The Netherlands; ²Guy's and St Thomas' NHS Foundation Trust, King's College London, London, UK; ³Clinical Division of Haematology and Haemostaseology, Department of Medicine I, Medical University of Vienna, Vienna, Austria; ⁴Department of General Internal Medicine, Cliniques Universitaires Saint Luc, UCL, Bruxelles, Belgium; ⁵Division of Angiology, Medical Clinic IV, University Hospital, Ludwig-Maximilians-University, Munich, Germany; ⁶National Pulmonary Hypertension Unit, Mater Misericordiae University Hospital, Dublin, Ireland; ⁷Respiratory Department, Ramón y Cajal Hospital, Madrid, Spain; ⁸Swiss Cardiovascular Centre, Division of Angiology, University of Bern, Bern University Hospital, Switzerland; ⁹Institute for Health Economics, Steinbeis-University, Berlin, Germany; ¹⁰LEGOS, Université Paris – Dauphine, Paris, France; ¹¹University of Castilla-La Mancha, Talavera de la Reina, Toledo, Spain; ¹²Department of Medicine, Université Paris Descartes, Paris, France; ¹³Institute for Pharmacology and Preventive Medicine, Berlin, Germany; ¹⁴Internal and Cardiovascular Medicine-Stroke Unit, University of Perugia, Perugia, Italy

Introduction

- Edoxaban, a direct FXa inhibitor, is approved for the treatment and secondary prevention of acute VTE, based on its comparable efficacy and superior safety versus warfarin during the 12 month Hokusai-VTE trial¹
- The ETNA-VTE-Europe (NCT02943993) study is being conducted to gain further insight into the efficacy and safety of extended treatment with edoxaban up to 18 months in routine clinical practice²
- There are data to suggest that the index clinical manifestation of venous thromboembolism (VTE; either deep vein thrombosis [DVT] or pulmonary embolism [PE] with or without DVT) strongly predicts the type of recurrence. However, data on the association between index manifestation and absolute recurrent VTE rates are conflicting³
- This poster assesses the patient characteristics and evaluates the rates and associations between the index VTE event (PE with or without DVT versus DVT alone) and 12-month outcomes


Methods

- ETNA-VTE-Europe is an 18-month, prospective, single-arm, non-interventional, multinational, post-authorisation safety study²
- A total of 2407 patients with acute symptomatic VTE who were receiving edoxaban in routine practice were enrolled from eight European Countries (Austria, Belgium, Germany, Ireland, Italy, the Netherlands, Switzerland and the United Kingdom)
- At 12 months, patients with an index event of PE with or without DVT (PE±DVT) were compared with those with an index event of DVT without PE

Results

- Patients with acute PE±DVT had a slightly higher age, a higher percentage of frailty, more hypertension and a more frequent history of PE than patients with DVT only (**Table 1**)
- Patients with DVT alone had more history of chronic venous insufficiency and an increased history of prior DVT (**Table 1**)
- Overall, VTE recurrence was 2.83%, rates of any bleeding 13.00%, any stroke 0.83%, major bleeding 1.91% and all-cause mortality 3.37% (**Figure 1–3**)
- Patients with index PE±DVT had numerically higher rates of all-cause mortality (4.08% versus 2.85%), cardiovascular mortality (1.69% versus 1.21%) and major bleeding (2.39% versus 1.57%) rates than those with index DVT alone (**Figure 2 and 3**)

Figure 1. Events of venous thromboembolism and stroke during 12-month follow-up




DVT, deep vein thrombosis; PE, pulmonary embolism; VTE, venous thromboembolism

Table 1. Patient baseline characteristics by type of index event

Patient characteristics	Overall [N=2407]	PE±DVT [N=1004]	DVT alone [N=1403]
Female, n (%)	1121 (46.6)	485 (48.3)	636 (45.3)
Age, years, mean ± SD	63.2 ± 15.87	64.4 ± 15.29	62.3 ± 16.23
Recalc. eGFR (Cockcroft-Gault, mL/min), mean ± SD	94.2 ± 38.97	92.8 ± 38.30	95.3 ± 39.47
Frailty (physician judgement), n (%)	306 (12.7)	140 (14.0)	166 (11.8)
Medical history, n (%)			
Hypertension	1041 (43.2)	476 (47.4)	565 (40.3)
Chronic venous insufficiency	264 (11.0)	75 (7.5)	189 (13.5)
COPD	162 (6.7)	86 (8.6)	76 (5.4)
Cancer	234 (9.7)	110 (11.0)	124 (8.8)
Cancer still active*	89 (38.2)	38 (34.9)	51 (41.1)
Bleeding history	71 (2.9)	47 (4.7)	24 (1.7)
History of major or CRNM bleeding	46 (1.9)	28 (2.8)	18 (1.3)
History of major bleeding	22 (0.9)	12 (1.2)	10 (0.7)
Ischaemic stroke	58 (2.4)	37 (3.7)	21 (1.5)
Prior VTE, n (%)			
Prior PE±DVT	177 (7.4)	112 (11.2)	65 (4.6)
Prior DVT only	388 (16.1)	110 (11.0)	278 (19.8)
Edoxaban treatment at baseline, n (%)			
Edoxaban 60 mg	2100 (87.2)	891 (88.7)	1209 (86.2)
Edoxaban 30 mg	307 (12.8)	113 (11.3)	194 (13.8)


*Percentage based on patients with cancer history
COPD, chronic obstructive pulmonary disease; CRNM, clinically relevant non-major; DVT, deep vein thrombosis; eGFR, estimated glomerular filtration rate; PE, pulmonary embolism; SD, standard deviation; VTE, venous thromboembolism

Figure 2. Bleeding events during 12-month follow-up



CRNM, clinically relevant non-major; DVT, deep vein thrombosis; GI, gastrointestinal; ICH, intracranial haemorrhage; ISTH, International Society on Thrombosis and Haemostasis; PE, pulmonary embolism

Figure 3. Deaths reported during 12-month follow-up



CV, cardiovascular; DVT, deep vein thrombosis; PE, pulmonary embolism

Conclusions

- Low rates of bleeding and venous thromboembolism events were reported after 12 months of treatment with edoxaban in clinical practice reinforcing the safety and effectiveness of edoxaban treatment in VTE patients
- Patients with index pulmonary embolism with or without deep vein thrombosis reported a higher mortality and bleeding events than those with deep vein thrombosis alone
- The 18-month outcomes results are awaited

References

- Büller HR, et al. *N Engl J Med*. 2013;369:1406-15
- Cohen AT, et al. *Thromb J*. 2018;16:9
- Mearns ES, et al. *J Thromb Haemost*. 2015;13:1043-52

Acknowledgements

This poster was sponsored by Daiichi Sankyo Europe GmbH, Munich, Germany. Writing and editorial support were provided by Sean Littlewood from inScience Communications, Springer Healthcare Ltd, UK and funded by Daiichi Sankyo Europe GmbH, Munich Germany

Declaration of interest

Michiel Coppens has received research funding and honoraria for consultancy or lecturing from Daiichi Sankyo, Boehringer Ingelheim, Bayer, Bristol-Myers Squibb, Portola, Sanquin Blood Supply, CSL Behring and Pfizer

